ELSEVIER

Journal of Organometallic Chemistry 544 (1997) 285–287

Journal of Organometallic Chemistry

Author Index of Volume 544

Agarkov, A.Yu., see Voskoboynikov, A.Z. 65
Alt, H.G., see Peifer, B. 115
Alt, H.G., see Schmid, C. 139
Alvarez, R., see Höpfl, H. 175
Arnason, I., see Björgvinsson, M. 207
Avtomonov, E.V., Megges, K., Li, X., Lorberth, J., Wocadlo, S., Massa, W., Harms, K., Churakov, A.V. and Howard, J.A.K., Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives, 79
Azam, K.A., Hossain, Md.M., Hursthouse, M.B., Kabir, S.E., Karim, M.M. and Malik, K.M.A., A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[$Mn_2(CO)_6\{\eta^1$-SC$(NMe_2)_2\}$], 23

Bajaj, H.C., see Das, A. 55
Beletskaya, I.P., see Voskoboynikov, A.Z. 65
Bhadbhade, M.M., see Das, A. 55
Björgvinsson, M., Halldorsson, S., Arnason, I., Magull, J. and Fenske, D., Preparation and characterization of $(C_5Me_4Ph)TiCl_3$, the oxochloride complexes $[(C_5Me_4Ph)TiCl_2]_2(\mu$-O) and $[(C_5Me_4Ph)TiCl(\mu$-O)$]_3$ and the oxo-complex $[(C_5Me_4Ph)Ti]_4(\mu$-O$)_6$. The X-ray crystal structures of $[(C_5Me_4Ph)TiCl_2]_2(\mu$-O) and $[(C_5Me_4Ph)Ti]_4(\mu$-O$)_6$, 207
Bosque, R., Caubet, A., López, C., Espinosa, E. and Molins, E., Syntheses, characterization and study of the properties of heterobimetallic compounds containing ferrocenyl units. X-ray crystal structure of: $[Zn\{(\eta^5$-$C_5H_5)Fe(\eta^5$-$C_5H_4)$–CH=N–CH_2–CH_2–$N(CH_3)_2\}]Cl_2$], 233
Bott, S.G., see Don, M.-J. 15
Bringmann, G., Stowasser, R. and Göbel, L., DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand, 7
Brüser, W., see Poremba, P. 1

Calhorda, M.J., see De Azevedo, C.G. 257
Carcelli, M., Delledonne, D., Fochi, A., Pelizzi, G., Rodríguez-Argüelles, M.C. and Russo, U., The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes, 29
Castillo, D., see Höpfl, H. 175
Caubet, A., see Bosque, R. 233
Cavell, K.J., see Desjardins, S.Y. 163
Cea-Olivares, R., see Flores-Santos, L. 37
Churakov, A.V., see Avtomonov, E.V. 79
Contreras, R., see Höpfl, H. 175
Corma, A., Iglesias, M., Mohino, F. and Sánchez, F., Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2*S*,4*R*)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2*S*,4*S*)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins, 147

Da Piedade, F.M., see De Azevedo, C.G. 257
Daran, J.-C., see Höpfl, H. 175
Das, A., Bajaj, H.C. and Bhadbhade, M.M., Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of $[C_5H_5FeC_5H_4CH=CHC_5H_4N$–$CH_3]PF_6$ with a mirror creating disorder, 55
De Azevedo, C.G., Calhorda, M.J., De C.T. Carrondo, M.A.A.F., Dias, A.R., Duarte, M.T., Galvão, A.M., Gamelas, C.A., Gonçalves, I.S., Da Piedade, F.M. and Romão, C.C., Nucleophilic and electrophilic reactions of C_5 *cyclo*-polyenes coordinated to the $[CpMoL_2]^{n+}$ fragment (n = 1,2; L = 1/2dppe, PMe_3, $P(OMe)_3$, CO), 257
De C.T. Carrondo, M.A.A.F., see De Azevedo, C.G. 257
Delledonne, D., see Carcelli, M. 29
Deravakian, A., see Melikyan, G.G. 143
Desjardins, S.Y., Cavell, K.J., Hoare, J.L., Skelton, B.W., Sobolev, A.N., White, A.H. and Keim, W., Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway, 163
Desmet, M., see Raubenheimer, H.G. 91
De Souza, A.-L.A.B., see Hussain, Z.I. 121
Dias, A.R., see De Azevedo, C.G. 257
Don, M.-J., Yang, K., Bott, S.G. and Richmond, M.G., Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex $[Cp^*Rh(mnt)]_2$. Confirmation of bridging and chelating mnt ligands in the solid-state structure, 15
Duarte, M.T., see De Azevedo, C.G. 257

Edelmann, F.T., see Poremba, P. 1
Espinosa, E., see Bosque, R. 233

Farfán, N., see Höpfl, H. 175
Fenske, D., see Björgvinsson, M. 207
Fernández, L., see Höpfl, H. 175
Fierro, J.L.G., Martínez-Ripoll, M., Merchán, M.D., Rodríguez, A., Terreros, P., Torrens, H. and Vivar-Cerrato, M.A., Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-$[Rh(SC_6F_5)(CO)(PPh_3)_2]$, 243
Filippou, A.C., Winter, J.G., Kociok-Köhn, G. and Hinz, I., Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes, 225
Flores-Santos, L., Cea-Olivares, R., Hernández-Ortega, S., Toscano, R.A., García-Montalvo, V., Novosad, J. and Woollins, J.D., First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se*,*Se'*) ligand. The synthesis and X-ray crystal structures of $Bu_2Sn[\{N(SePPh_2)_2$-*Se*,*Se'*$\}_2]$ and Ph_2Cl-$Sn[N(SePPh_2)_2$-*Se*,*Se'*$]\cdot H_2O$37
Fochi, A., see Carcelli, M. 29

Galván, M., see Höpfl, H. 175
Galvão, A.M., see De Azevedo, C.G. 257
Gamelas, C.A., see De Azevedo, C.G. 257
García-Montalvo, V., see Flores-Santos, L. 37
Göbel, L., see Bringmann, G. 7
Gonçalves, I.S., see De Azevedo, C.G. 257
Gutmann, T., see Schenk, W.A. 69

Elsevier Science S.A.

Halldorsson, S., see Björgvinsson, M. 207
Halut, S., see Höpfl, H. 175
Harms, K., see Avtomonov, E.V. 79
Hernández-Ortega, S., see Flores-Santos, L. 37
Herzog, U. and Roewer, G., Preparation of oligosilanes containing perhalogenated silyl groups ($-SiX_3$, $-SiX_2-$, $>SiX-$, X = Cl, Br) and their hydrogenation by stannanes, 217
Hinz, I., see Filippou, A.C. 225
Hoare, J.L., see Desjardins, S.Y. 163
Hockless, D.C.R., see Whittall, I.R. 189
Hockless, D.C.R., see Whittall, I.R. 277
Höpfl, H., Farfán, N., Castillo, D., Santillan, R., Contreras, R., Martínez-Martínez, F.J., Galván, M., Alvarez, R., Fernández, L., Halut, S. and Daran, J.-C., Dynamic NMR and X-ray diffraction study of (N–B)-diphenyl(2-aminoethoxy)borane derivatives of ephedrines and pseudoephedrines, 175
Hossain, Md.M., see Azam, K.A. 23
Houbrechts, S., see Whittall, I.R. 277
Howard, J.A.K., see Avtomonov, E.V. 79
Howard, J.A.K., see Voskoboynikov, A.Z. 65
Hrušak, J., see Raubenheimer, H.G. 91
Humphrey, M.G., see Whittall, I.R. 189
Humphrey, M.G., see Whittall, I.R. 277
Hursthouse, M.B., see Azam, K.A. 23
Hussain, Z.I., De Souza, A.-L.A.B. and Whiteley, M.W., Investigations on the synthesis and stability of η^3-bonded cycloheptadienyl complexes of molybdenum, 121

Iglesias, M., see Corma, A. 147
Ishikawa, M., see Ohshita, J. 49

Johnsen, E., see Roberts, R.M.G. 197
Jones, R.G., see Wiseman, A.J. 129

Kabir, S.E., see Azam, K.A. 23
Karim, M.M., see Azam, K.A. 23
Keim, W., see Desjardins, S.Y. 163
Kociok-Köhn, G., see Filippou, A.C. 225
Kruger, G.J., see Raubenheimer, H.G. 91
Kuz'mina, L.O., see Voskoboynikov, A.Z. 65

Leung, S.-W., see Singleton, D.A. 157
Li, X., see Avtomonov, E.V. 79
Lindeque, L., see Raubenheimer, H.G. 91
López, C., see Bosque, R. 233
Lorberth, J., see Avtomonov, E.V. 79
Luther-Davies, B., see Whittall, I.R. 189

Maes, J., see Whittall, I.R. 277
Magull, J., see Björgvinsson, M. 207
Malik, K.M.A., see Azam, K.A. 23
Martínez-Martínez, F.J., see Höpfl, H. 175
Martínez-Ripoll, M., see Fierro, J.L.G. 243
Masaoka, S., see Ohshita, J. 49
Massa, W., see Avtomonov, E.V. 79
Megges, K., see Avtomonov, E.V. 79
Melikyan, G.G. and Deravakian, A., Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol, 143
Merchán, M.D., see Fierro, J.L.G. 243
Milius, W., see Schmid, C. 139
Mohino, F., see Corma, A. 147
Molins, E., see Bosque, R. 233

Nekoda, E., see Ohshita, J. 49
Noltemeyer, M., see Poremba, P. 1
Novosad, J., see Flores-Santos, L. 37

Ohshita, J., Nekoda, E., Masaoka, S. and Ishikawa, M., Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents, 49
Oldenburg, K. and Vogler, A., Optical metal-to-metal charge transfer in $[Fe^{II}(C_5H_4PPh_2)_2Re^{I}(CO)_3Cl]^+$, 101
Olivier, P.J., see Raubenheimer, H.G. 91

Peifer, B., Welch, M.B. and Alt, H.G., Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation, 115
Pelizzi, G., see Carcelli, M. 29
Persoons, A., see Whittall, I.R. 277
Plenio, H. and Warnecke, A., η^5-Complexes of cyclopentadienylsilylethers ($C_5H_4OSiR_3$) and hydroxycyclopentadiene (C_5H_4OH) with titanium and zirconium chlorides, 133
Poremba, P., Reißmann, U., Noltemeyer, M., Schmidt, H.-G., Brüser, W. and Edelmann, F.T., Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands, 1

Raubenheimer, H.G., Olivier, P.J., Lindeque, L., Desmet, M., Hrušak, J. and Kruger, G.J., Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds, 91
Reißmann, U., see Poremba, P. 1
Richmond, M.G., see Don, M.-J. 15
Roberts, R.M.G. and Johnsen, E., Synthesis, stereochemistry and decomplexation of (η-arene)(η-cyclopentadienyl) iron(II) hexafluorophosphate complexes containing amino acid side chains. A route to *N*-arylamino acids, 197
Rodríguez, A., see Fierro, J.L.G. 243
Rodríguez-Argüelles, M.C., see Carcelli, M. 29
Roewer, G., see Herzog, U. 217
Romão, C.C., see De Azevedo, C.G. 257
Russo, U., see Carcelli, M. 29

Samoc, M., see Whittall, I.R. 189
Sánchez, F., see Corma, A. 147
Santillan, R., see Höpfl, H. 175
Schenk, W.A. and Gutmann, T., Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes, 69
Schmid, C., Alt, H.G. and Milius, W., Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums, 139
Schmid, S., see Whittall, I.R. 277
Schmidt, H.-G., see Poremba, P. 1
Sebald, A., see Wrackmeyer, B. 105
Shestakova, A.K., see Voskoboynikov, A.Z. 65
Singleton, D.A. and Leung, S.-W., In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes, 157
Skelton, B.W., see Desjardins, S.Y. 163
Sobolev, A.N., see Desjardins, S.Y. 163
Stowasser, R., see Bringmann, G. 7

Terreros, P., see Fierro, J.L.G. 243
Tian, G., see Zhang, Y. 43
Torrens, H., see Fierro, J.L.G. 243
Toscano, R.A., see Flores-Santos, L. 37

Vivar-Cerrato, M.A., see Fierro, J.L.G. 243
Vogler, A., see Oldenburg, K. 101
Voskoboynikov, A.Z., Agarkov, A.Yu., Shestakova, A.K., Beletskaya, I.P., Kuz'mina, L.O. and Howard, J.A.K., Synthesis and molecular structure of $Na[[1,3\text{-}(Me_3Si)_2\text{-}C_5H_3]Yb^{II}]_2(\mu\text{-I})$, 65

Warnecke, A., see Plenio, H. 133
Welch, M.B., see Peifer, B. 115
Went, M.J., see Wiseman, A.J. 129
White, A.H., see Desjardins, S.Y. 163
Whiteley, M.W., see Hussain, Z.I. 121
Whittall, I.R., Humphrey, M.G., Houbrechts, S., Maes, J., Persoons, A., Schmid, S. and Hockless, D.C.R., Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of $1\text{-}(HC\equiv C)\text{-}3,5\text{-}C_6H_3$ (*trans*-$C\equiv CRuCl(dppm)_2)_2$ and $1,3,5\text{-}C_6H_3(C\equiv CAu(PPh_3))_3$, 277
Whittall, I.R., Humphrey, M.G., Samoc, M., Luther-Davies, B. and Hockless, D.C.R., Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of $Au(C\equiv CPh)(nmdpp)$ and $Au((E)\text{-}4,4'\text{-}C\equiv CC_6H_4CH=CHC_6H_4\text{-}NO_2)(nmdpp)$, 189
Winter, J.G., see Filippou, A.C. 225
Wiseman, A.J., Jones, R.G. and Went, M.J., Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane), 129
Wocadlo, S., see Avtomonov, E.V. 79
Woollins, J.D., see Flores-Santos, L. 37

Wrackmeyer, B. and Sebald, A., Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes, 105

Xu, S., see Zhang, Y. 43

Yang, K., see Don, M.-J. 15

Zhang, W., see Zhang, Y. 43

Zhang, Y., Xu, S., Tian, G., Zhang, W. and Zhou, X., Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes $(Me_2E)[(\eta^5\text{-}C_5H_4)M(CO)]_2(\mu\text{-}CO)_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of $(Me_2Ge)[(\eta^5\text{-}C_5H_4)Fe(CO)_2]_2SnCl_2$, 43

Zhou, X., see Zhang, Y. 43

ELSEVIER

Journal of Organometallic Chemistry 544 (1997) 289–294

Journal of Organometallic Chemistry

Subject Index of Volume 544

Acetylide
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C_6H_3(*trans*-C≡CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C≡CAu(PPh_3))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4′-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
Alkynes
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
Alkynyl
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4′-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC≡C)-3,5-C_6H_3(*trans*-C≡CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C≡CAu(PPh_3))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
Allyl
Investigations on the synthesis and stability of η^3-bonded cycloheptadienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121
Arenes
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129
Arsenic
Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79
Arsenic amides
Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79
Arylnickel complexes
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163
Biaryl lactones
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
Bimetallic complexes
Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69
Bis(silyl) ketone
Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49
Boron
In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105
Bridged complexes
Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes $(Me_2E)[(\eta^5\text{-}C_5H_4)M(CO)]_2(\mu\text{-}CO)_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of $(Me_2Ge)[(\eta^5\text{-}C_5H_4)Fe(CO)_2]_2SnCl_2$ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43
Bridging ligand
Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex $[Cp^*Rh(mnt)]_2$. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15
Carbene complexes
Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušak and G.J. Kruger), 91
Carbon monoxide
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163
Carbonohydrazone
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorgan-

Elsevier Science S.A.

otin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29

Carbonyl
A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[$Mn_2(CO)_9\{\eta^1$-SC(NMe$_2$)$_2$}] (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23

Charge transfer
Optical metal-to-metal charge transfer in [$Fe^{III}(C_5H_4PPh_2)_2$-$Re^I(CO)_3Cl]^+$ (K. Oldenburg and A. Vogler), 101

Chelating ligand
Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [$Cp^*Rh(mnt)]_2$. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15

Chiral
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4'-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Chromiumtricarbonyl rotor
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

C_5Me_4Ph ligand
Preparation and characterization of (C_5Me_4Ph)$TiCl_3$, the oxochloride complexes [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)TiCl(μ-O)]$_3$ and the oxo-complex [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$. The X-ray crystal structures of [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

Copolymerisation
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Correlated dynamics
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

Cycloheptadienyl
Investigations on the synthesis and stability of η^3-bonded cycloheptadienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121

Cyclooctatetraenyl ligands
Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1

Cyclopentadienyl arsenic compounds: crystal structure
Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79

Cyclopentadienylsilylether
η^5-Complexes of cyclopentadienylsilylethers ($C_5H_4OSiR_3$) and hydroxycyclopentadiene (C_5H_4OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133

Density-functional calculation
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

Diastereoselectivity
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143

Diels–Alder reactions
In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157

Diorganotin derivative
First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se,Se'*) ligand. The synthesis and X-ray crystal structures of Bu_2Sn[{$N(SePPh_2)_2$-*Se,Se'*}$_2$] and Ph_2ClSn[$N(SePPh_2)_2$-*Se,Se'*]·H_2O (L. Flores-Santos, R. Cea-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37

Ethene
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Ferrocene
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [$C_5H_5FeC_5H_4CH$ = CHC_5H_4*N*–CH_3]PF_6 with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

Ferrocenium
Optical metal-to-metal charge transfer in [$Fe^{III}(C_5H_4PPh_2)_2$-$Re^I(CO)_3Cl]^+$ (K. Oldenburg and A. Vogler), 101

Fluorenyl
Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Fulvene
Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Germanium
Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhn and I. Hinz), 225

Germylene
Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhn and I. Hinz), 225

Gold
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC ≡ C)-3,5-C_6H_3(*trans*-C ≡ CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C ≡ CAu(PPh_3))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4′-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Gold (I) and gold (III) compounds

Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušak and G.J. Kruger), 91

Group 15

Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79

Hafnium

Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139

Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Hapticity

Investigations on the synthesis and stability of η^3-bonded cycloheptadienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121

Hydroformylation

Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[Rh(SC_6F_5)(CO)PPh_3)$_2$] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243

Hydrogenation

Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147

Hydroxycyclopentadiene

η^5-Complexes of cyclopentadienylsilylethers ($C_5H_4OSiR_3$) and hydroxycyclopentadiene (C_5H_4OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133

Imidobis ligand

First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se*,*Se*′) ligand. The synthesis and X-ray crystal structures of Bu_2Sn[{N(SePPh$_2$)$_2$-*Se*,*Se*′}$_2$] and Ph_2ClSn[N(SePPh$_2$)$_2$-*Se*,*Se*′]·H_2O (L. Flores-Santos, R. Cea-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37

Imino arsanes

Syntheses and structures of cyclopentadienyl arsenic compounds. II. Pentamethyl- and tetraisopropylcyclopentadienyl arsenic amido derivatives (E.V. Avtomonov, K. Megges, X. Li, J. Lorberth, S. Wocadlo, W. Massa, K. Harms, A.V. Churakov and J.A.K. Howard), 79

Indenyl

Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139

Insertion

Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhn and I. Hinz), 225

Insertion reaction

Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes (Me_2E)(η^5-C_5H_4)M(CO)]$_2$(μ-CO)$_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me_2Ge)(η^5-C_5H_4)Fe(CO)$_2$]$_2SnCl_2$ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43

Iron

Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes (Me_2E)(η^5-C_5H_4)M(CO)]$_2$(μ-CO)$_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me_2Ge)(η^5-C_5H_4)Fe(CO)$_2$]$_2SnCl_2$ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43

Kinetics

Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [$C_5H_5FeC_5H_4CH$ = CHC_5H_4N–CH_3]PF_6 with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

Kurtz

Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4′-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189

Ligand coupling

Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

Manganese

A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[$Mn_2(CO)_8${η^1-SC(NMe_2)$_2$}] (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23

Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Metal carbonyl

Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes (Me_2E)(η^5-C_5H_4)M(CO)]$_2$(μ-CO)$_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of (Me_2Ge)(η^5-C_5H_4)Fe(CO)$_2$]$_2SnCl_2$ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43

Molybdenum

Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

Insertion of $GeCl_2$ into molybdenum–hydrogen bonds: a convenient route to dichlorogermyl complexes (A.C. Filippou, J.G. Winter, G. Kociok-Köhn and I. Hinz), 225

Investigations on the synthesis and stability of η^3-bonded cycloheptadienyl complexes of molybdenum (Z.I. Hussain, A.-L.A.B. De Souza and M.W. Whiteley), 121

Nickel

Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal struc-

tures of 1-(HC ≡ C)-3,5-C_6H_3(*trans*-C ≡ CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C ≡ CAu(PPh_3))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

NMR
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Nonlinear optics
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC ≡ C)-3,5-C_6H_3(*trans*-C ≡ CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C ≡ CAu(PPh_3))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277

Olefin polymerization
Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Oligosilanes
Preparation of oligosilanes containing perhalogenated silyl groups (–SiX_3, –SiX_2–, >SiX–, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217

Organoboration
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Organolanthanide complexes
Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1

Organolithium reagent
Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49

Organotin compound
First functionalized diorganotin (IV) derivative containing the imidobis(diphenylselenophosphinate-*Se*,*Se'*) ligand. The synthesis and X-ray crystal structures of $Bu_2Sn[\{N(SePPh_2)_2$-*Se*,*Se'*$\}_2]$ and $Ph_2ClSn[N(SePPh_2)_2$-*Se*,*Se'*$]\cdot H_2O$ (L. Flores-Santos, R. Cea-Olivares, S. Hernández-Ortega, R.A. Toscano, V. García-Montalvo, J. Novosad and J.D. Woollins), 37

Organotin compounds
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29

Oxochloride complexes
Preparation and characterization of (C_5Me_4Ph)$TiCl_3$, the oxochloride complexes [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)TiCl(μ-O)]$_3$ and the oxo-complex [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$. The X-ray crystal structures of [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

Pentamethylcylopentadienyl rhodium complex
Redox investigation and X-ray diffraction structure of the dimeric rhodium maleonitriledithiolate complex [Cp* Rh(mnt)]$_2$. Confirmation of bridging and chelating mnt ligands in the solid-state structure (M.-J. Don, K. Yang, S.G. Bott and M.G. Richmond), 15

Perhalogenated silyl groups
Preparation of oligosilanes containing perhalogenated silyl groups (–SiX_3, –SiX_2–, >SiX–, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217

Phenyltetramethylcyclopentadiene
Preparation and characterization of (C_5Me_4Ph)$TiCl_3$, the oxochloride complexes [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)TiCl(μ-O)]$_3$ and the oxo-complex [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$. The X-ray crystal structures of [(C_5Me_4Ph)$TiCl_2$]$_2$(μ-O) and [(C_5Me_4Ph)Ti]$_4$(μ-O)$_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207

Phosphorus
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Physicochemical studies
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [$C_5H_5FeC_5H_4CH$ = CHC_5H_4N–CH_3]PF_6 with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

Platinum
Reactivity of 1-alkynylplatinum(II) complexes towards trialkylboranes (B. Wrackmeyer and A. Sebald), 105

Polydentate ligands
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29

Polymerisation
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Polynuclear complexes
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of [$C_5H_5FeC_5H_4CH$ = CHC_5H_4N–CH_3]PF_6 with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

Polysilanecarboxylate
Reactions of tris(trimethylsilyl)silanecarboxylates with organolithium reagents (J. Ohshita, E. Nekoda, S. Masaoka and M. Ishikawa), 49

Polysilanes
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Propargyl cation
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143

Propargyl radical
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143

Reductive dimerization
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143

Regioisomers
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7

Rhenium carbonyl complex
Optical metal-to-metal charge transfer in [$Fe^{III}(C_5H_4PPh_2)_2$-$Re^I(CO)_3Cl]^+$ (K. Oldenburg and A. Vogler), 101

Rhodium
Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147
Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-$[Rh(SC_6F_5)(CO)(PPh_3)_2]$ (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243
Ruthenium
Insertion of $SnCl_2$ into the metal–metal bonds in bridged binuclear complexes $(Me_2E)(\eta^5\text{-}C_5H_4)M(CO)]_2(\mu\text{-}CO)_2$ (E = Si, Ge; M = Fe, Ru). The molecular structure of $(Me_2Ge)(\eta^5\text{-}C_5H_4)Fe(CO)_2]_2SnCl_2$ (Y. Zhang, S. Xu, G. Tian, W. Zhang and X. Zhou), 43
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC ≡ C)-3,5-C_6H_3(*trans*-C ≡ CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C ≡ CAu(PPh$_3$))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129

Sandwich complexes
Cyclooctatetraenyl complexes of the early transition metals and lanthanides VIII. Anionic sandwich complexes of the lanthanides containing silylated cyclooctatetraenyl ligands (P. Poremba, U. Reißmann, M. Noltemeyer, H.-G. Schmidt, W. Brüser and F.T. Edelmann), 1
Semicarbazone
The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds. Part I: the diorganotin(IV) complexes (M. Carcelli, D. Delledonne, A. Fochi, G. Pelizzi, M.C. Rodríguez-Argüelles and U. Russo), 29
Silanes
Synthesis of manganese and ruthenium complexes of phenyltrimethylsilane, diphenyltetramethyldisilane and poly(methylphenylsilane) (A.J. Wiseman, R.G. Jones and M.J. Went), 129
Silicon
In situ formation of alkenyl- and alkynylboranes for Diels–Alder reactions by boron–silicon exchange with alkenyl- and alkynylsilanes (D.A. Singleton and S.-W. Leung), 157
Preparation of oligosilanes containing perhalogenated silyl groups (–SiX_3, –SiX_2–, >SiX–, X = Cl, Br) and their hydrogenation by stannanes (U. Herzog and G. Roewer), 217
Synthesis and molecular structure of $Na\{[1,3\text{-}(Me_3Si)_2C_5H_3]\text{-}Yb^{II}\}_2(\mu\text{-}I)$ (A.Z. Voskoboynikov, A.Yu. Agarkov, A.K. Shestakova, I.P. Beletskaya, L.O. Kuz'mina and J.A.K. Howard), 65
Single-electron transfer
Tetrahydrofuran as an one-electron donor: Highly diastereoselective coupling of cobalt-complexed propargyl alcohol (G.G. Melikyan and A. Deravakian), 143
Sterically crowded
Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139
Steric and electronic effects
DF-studies on a 'stop and go' rotor: steric and electronic factors determining the regio- and stereochemical position of a η^6-$Cr(CO)_3$ metal fragment on a helically distorted biaryl ligand (G. Bringmann, R. Stowasser and L. Göbel), 7
Supported catalysts
Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147
Susceptibility
Organometallic complexes for nonlinear optics. 14. Syntheses and second-order nonlinear optical properties of ruthenium, nickel and gold σ-acetylides of 1,3,5-triethynylbenzene: X-ray crystal structures of 1-(HC ≡ C)-3,5-C_6H_3(*trans*-C ≡ CRuCl(dppm)$_2$)$_2$ and 1,3,5-C_6H_3(C ≡ CAu(PPh$_3$))$_3$ (I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless), 277
Organometallic complexes for non-linear optics XII. Syntheses and second-order susceptibilities of (neomenthyldiphenylphosphine)gold σ-arylacetylides: X-ray crystal structures of Au(C≡CPh)(nmdpp) and Au((*E*)-4,4'-C≡CC_6H_4CH=CHC_6H_4-NO_2)(nmdpp) (I.R. Whittall, M.G. Humphrey, M. Samoc, B. Luther-Davies and D.C.R. Hockless), 189
Synthesis
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of $[C_5H_5FeC_5H_4CH = CHC_5H_4N\text{–}CH_3]PF_6$ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

Thiazolyl derivatives
Oxidative addition of mono and bis(carbene) complexes derived from imidazolyl and thiazolyl gold(I) compounds (H.G. Raubenheimer, P.J. Olivier, L. Lindeque, M. Desmet, J. Hrušak and G.J. Kruger), 91
Thiolate
Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-$[Rh(SC_6F_5)(CO)(PPh_3)_2]$ (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243
Thiourea
A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-$[Mn_2(CO)_9\{\eta^1\text{-}SC(NMe_2)_2\}]$ (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23
Ti_4O_6 cage
Preparation and characterization of $(C_5Me_4Ph)TiCl_3$, the oxochloride complexes $[(C_5Me_4Ph)TiCl_2]_2(\mu\text{-}O)$ and $[(C_5Me_4Ph)TiCl(\mu\text{-}O)]_3$ and the oxo-complex $[(C_5Me_4Ph)Ti]_4(\mu\text{-}O)_6$. The X-ray crystal structures of $[(C_5Me_4Ph)TiCl_2]_2(\mu\text{-}O)$ and $[(C_5Me_4Ph)Ti]_4(\mu\text{-}O)_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
(η^5-cyclopentadienyl)titanium
Preparation and characterization of $(C_5Me_4Ph)TiCl_3$, the oxochloride complexes $[(C_5Me_4Ph)TiCl_2]_2(\mu\text{-}O)$ and $[(C_5Me_4Ph)TiCl(\mu\text{-}O)]_3$ and the oxo-complex $[(C_5Me_4Ph)Ti]_4(\mu\text{-}O)_6$. The X-ray crystal structures of $[(C_5Me_4Ph)TiCl_2]_2(\mu\text{-}O)$ and $[(C_5Me_4Ph)Ti]_4(\mu\text{-}O)_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
Titanium chloride
η^5-Complexes of cyclopentadienylsilylethers ($C_5H_4OSiR_3$) and hydroxycyclopentadiene (C_5H_4OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133

Tungsten
Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69

X-ray diffraction
A dinuclear manganese carbonyl complex of tetramethylthiourea: X-ray crystal structure of *eq*-[$Mn_2(CO)_8\{\eta^1$-$SC(NMe_2)_2\}$] (K.A. Azam, Md.M. Hossain, M.B. Hursthouse, S.E. Kabir, M.M. Karim and K.M.A. Malik), 23
Rhodium(I) fluorothiolate complexes as hydroformylation catalyst precursors. Crystal structure of two polymorphs of *trans*-[$Rh(SC_6F_5)(CO)(PPh_3)_2$] (J.L.G. Fierro, M. Martínez-Ripoll, M.D. Merchán, A. Rodríguez, P. Terreros, H. Torrens and M.A. Vivar-Cerrato), 243

X-ray structure
Preparation and characterization of $(C_5Me_4Ph)TiCl_3$, the oxochloride complexes $[(C_5Me_4Ph)TiCl_2]_2(\mu$-O) and $[(C_5Me_4Ph)TiCl(\mu$-$O)]_3$ and the oxo-complex $[(C_5Me_4Ph)Ti]_4(\mu$-$O)_6$. The X-ray crystal structures of $[(C_5Me_4Ph)TiCl_2]_2(\mu$-O) and $[(C_5Me_4Ph)Ti]_4(\mu$-$O)_6$ (M. Björgvinsson, S. Halldorsson, I. Arnason, J. Magull and D. Fenske), 207
Redox responsive bi/tri-nuclear complexes incorporating ferrocenyl unit: Synthesis, characterization, physicochemical studies and X-ray structure of $[C_5H_5FeC_5H_4CH=CHC_5H_4N$–$CH_3]PF_6$ with a mirror creating disorder (A. Das, H.C. Bajaj and M.M. Bhadbhade), 55

X-ray structures
Single component N–O chelated arylnickel(II) complexes as ethene polymerisation and CO/ethene copolymerisation catalysts. Examples of ligand induced changes to the reaction pathway (S.Y. Desjardins, K.J. Cavell, J.L. Hoare, B.W. Skelton, A.N. Sobolev, A.H. White and W. Keim), 163

Ytterbium
Synthesis and molecular structure of $Na\{[1.3\text{-}(Me_3Si)_2C_5H_3]\text{-}Yb^{II}\}_2(\mu$-I) (A.Z. Voskoboynikov, A.Yu. Agarkov, A.K. Shestakova, I.P. Beletskaya, L.O. Kuz'mina and J.A.K. Howard), 65

Zeolites
Heterogenised catalysts on zeolites. Synthesis of new chiral Rh(I) complexes with (2S,4R)-*trans*-4-RCOO-2-(*t*-butylaminocarbonyl)pyrrolidines and (2S,4S)-*cis*-4-RCONH-2-(*t*-butylaminocarbonyl)pyrrolidines. Heterogenisation on silica and a USY-zeolite and study of the role of support on their catalytic profile in hydrogenation of olefins (A. Corma, M. Iglesias, F. Mohino and F. Sánchez), 147

Zirconium
Bimetallic complexes III. Synthesis and reactions of $C_5H_4PPh_2$-bridged zirconium–molybdenum and zirconium–tungsten complexes (W.A. Schenk and T. Gutmann), 69
Sterisch überladene Indenylkomplexe des Zirconiums und Hafniums (C. Schmid, H.G. Alt and W. Milius), 139
Synthese und Charakterisierung von C_1- und C_2-verbrückten Bis(fluorenyl)komplexen des Zirconiums und Hafniums und deren Anwendung bei der katalytischen Olefinpolymerisation (B. Peifer, M.B. Welch and H.G. Alt), 115

Zirconium chloride
η^5-Complexes of cyclopentadienylsilylethers ($C_5H_4OSiR_3$) and hydroxycyclopentadiene (C_5H_4OH) with titanium and zirconium chlorides (H. Plenio and A. Warnecke), 133